SECTION 18 EMERGENCY EXEMPTION FOR USE AND DISTRIBUTION ONLY WITHIN THE STATE OF NORTH CAROLINA FOR CONTROL OF BROWN MARMORATED STINK BUGS IN POME AND STONE FRUIT.

NORTH CAROLINA: <u>STATE RESTRICTED USE PESTICIDE</u> WHEN USED IN NORTH CAROLINA, APPLICATIONS CAN ONLY BE MADE BY LICENSED/CERTIFIED APPLICATORS OR BY PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE LICENSED/CERTIFIED APPLICATORS CERTIFICATION.



- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, Worker Protection Standard (WPS) requirements, and precautions on the federally registered product label for Scorpion 35 SL Insecticide (EPA Reg. No. 10163-317), as well as the Section 18 directions, must be followed.
- The additional spray drift recommendations on the registered product label under "Air Assisted Tree and Vine Sprayers (Berry / Small Fruit and Tuberous / Corm Vegetables only)" must also be followed.
- This label must be in the possession of the user at the time of pesticide application.
- Do not use Scorpion 35 SL Insecticide on pome and stone fruit after the expiration date of 10-15-2020.



Application restrictions exist for this product because of risk to bees and other insect pollinators. Follow application restrictions found in the directions for use for protection of pollinators on the registered product label for Scorpion® 35 SL Insecticide (EPA Reg. No. 10163-317). Do not apply this product while bees are foraging. This compound is highly toxic to honey bees. The persistence of residues and potential residual toxicity of dinotefuran in nectar and pollen suggests the possibility of chronic toxic risk to honey bee larvae and the eventual instability of the hive. This product is toxic to bees exposed to residues for more than 38 hours following treatment.

DIRECTIONS FOR USE

Direction for use on Pome and Stone Fruits			
PEST	RATE FL. OZ./A	REMARKS	RESTRICTIONS
Brown Marmorated Stink Bug	8 - 12 fl oz/A (0.203 lbs - 0.304 lbs ai/A)	complete coverage. Higher water volumes may provide improved insect control. Begin application when field sampling indicates pest activity. Use higher label rates when pest pressure increases and especially when adults are actively migrating into the orchard. Repeat as needed to maintain control,	 Do not apply this product until after petal fall. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Foliar Application Only Apply with ground equipment in adequate water to obtain uniform coverage (For best results, use at least 50 gallons/acre for ground applications).

Scorpion® is a registered trademark of Gowan Company, LLC.



Gowan Company P.O. BOX 5569 Yuma, AZ 85366-5569